

Review Article

Biological control of *Salmonella* on egg surfaces using plant-based compounds: A review

Nurten Yılmaz^{1,2}

¹Department of Medicinal and Aromatic Plants, Vocational School of Karaisali, Çukurova University, Adana, Türkiye;

²Biotechnology Research and Application Center, Çukurova University, Adana, Türkiye

Abstract

Salmonella contamination of shell eggs is a major public health concern worldwide and continues to challenge the poultry industry. Although conventional chemical sanitizers are effective, concerns regarding chemical residues, environmental impact, and consumer demand for clean-label products have accelerated interest in plant-derived antimicrobials. This review summarizes recent advances in the application of essential oils and plant extracts for controlling *Salmonella* on eggshell surfaces. Particular attention is given to the antimicrobial mechanisms of major bioactive compounds, including thymol, carvacrol, eugenol, cinnamaldehyde, and phenolic-rich plant extracts. The review also discusses practical application strategies, including washing, spraying, nanoemulsion-based delivery, and active packaging, highlighting their effectiveness in enhancing antimicrobial stability and prolonging egg shelf life. Finally, current challenges related to formulation standardization, industrial scalability, and regulatory approval are addressed, together with future perspectives for integrating plant-derived antimicrobials into sustainable egg sanitation systems.

Keywords: Active packaging, nanoemulsion, natural antimicrobials, plant extracts, *Salmonella enterica*

Introduction

Despite their exceptional nutritional value, eggs remain highly susceptible to microbial contamination during production, handling, transportation, and storage. Among the diverse foodborne pathogens associated with eggs, *Salmonella enterica*, particularly *Salmonella* Enteritidis, represents one of the most significant public health concerns because of its remarkable ability to survive on eggshell surfaces, penetrate the shell structure, and ultimately contaminate internal egg contents (Illera et al., 2022; Oliveira et al., 2022).

Consequently, effective decontamination of eggshell surfaces has become a critical prerequisite for reducing foodborne outbreaks, improving consumer safety, and extending the commercial shelf life of table eggs. Although synthetic sanitizers are capable of reducing microbial loads under industrial conditions, their widespread use has generated increasing concern regarding chemical residues, occupational exposure, environmental contamination, deterioration of eggshell quality, and the possible selection of tolerant bacterial populations after repeated application.

* **Correspondence:** N. Yılmaz (ntoy@cu.edu.tr).

<https://doi.org/10.30466/fsp.2026.57392.1035>

Received: 18 July 2026 **Accepted:** 04 August 2026

© 2026 Urmia University. All rights reserved.

Available online: 15 August 2026

Formaldehyde fumigation, for example, remains one of the most effective sanitization approaches for hatching eggs but has been increasingly criticized because of its toxicity toward both humans and developing embryos, motivating intensive research into safer alternatives (El-kashef & El Sabry, 2025; El-Sabroun et al., 2026).

Growing consumer preference for minimally processed foods and clean-label preservation strategies has substantially accelerated the development of naturally derived antimicrobial compounds for food protection. Instead of relying solely on synthetic disinfectants, contemporary food preservation increasingly exploits bioactive compounds extracted from plants, microorganisms, and renewable biological materials (Alkan & Yemenicioğlu, 2016). Essential oils (EOs), bacteriocins, edible biopolymers, plant phenolics, organic acids, and antimicrobial peptides have attracted considerable attention because they combine broad-spectrum antimicrobial activity with relatively favorable toxicological profiles and compatibility with sustainable food production systems (Karnwal & Malik, 2024; Teshome et al., 2022).

Natural antimicrobial compounds exhibit several advantages over conventional chemical sanitizers. Rather than acting through a single cellular target, many phytochemicals simultaneously disrupt bacterial membranes, interfere with ATP generation, alter membrane permeability, induce oxidative stress, inhibit quorum sensing, and suppress biofilm formation (Zhang et al., 2026). Such multi-target antimicrobial mechanisms considerably reduce the probability of bacterial adaptation while providing broad antimicrobial spectra against Gram-positive and Gram-negative pathogens, including *Salmonella*, *Escherichia coli*, *Listeria monocytogenes*, and *Staphylococcus aureus* (Raikwar et al., 2024). Among naturally occurring antimicrobials, EOs have emerged as particularly promising candidates for eggshell sanitation. Phenolic constituents such as thymol, carvacrol, eugenol, cinnamaldehyde, limonene, citral, and allyl isothiocyanate exhibit potent bactericidal activity by rapidly penetrating bacterial membranes, disrupting phospholipid

bilayers, increasing membrane permeability, and causing leakage of intracellular constituents (Allen et al., 2023; Upadhyaya et al., 2013).

Several experimental investigations have demonstrated that plant-derived antimicrobials can achieve remarkable reductions in *Salmonella* contamination within extremely short exposure periods (Elkomy et al., 2023; Upadhyaya et al., 2016). Despite these considerable advances, several technological challenges remain before natural antimicrobial interventions can be fully integrated into commercial egg production. Variability in EOs composition, limited stability of volatile bioactive compounds, interactions between antimicrobials and coating matrices, possible alterations in sensory quality, regulatory approval, industrial scalability, and economic feasibility continue to require comprehensive investigation (Konfo et al., 2023; Y. Zhang et al., 2026). Nevertheless, continuous progress in formulation technologies, nano-delivery systems, smart active packaging, and multidisciplinary preservation strategies strongly suggests that natural antimicrobial compounds will become integral components of future egg sanitation systems. Accordingly, this review critically summarizes current knowledge regarding the biological control of *Salmonella* on eggshell surfaces using natural antimicrobial compounds, emphasizing antimicrobial mechanisms, application technologies, coating systems, operational limitations, and future perspectives for sustainable egg safety.

Mechanisms of *Salmonella* contamination in eggs

Eggshell contamination primarily occurs after oviposition when freshly laid eggs are exposed to fecal material, litter, dust, contaminated nesting environments, processing equipment, and human handling. During this stage, *S. Enteritidis* readily colonizes the shell surface, where its subsequent fate depends largely on environmental conditions including temperature, relative humidity, storage duration, and the integrity of the eggshell cuticle (El-Sabroun et al., 2026; Oliveira et al., 2022). Initial bacterial attachment to the shell surface involves reversible physicochemical interactions that develop

into irreversible adhesion through exopolysaccharides, fimbriae, and surface proteins. Subsequent biofilm formation enhances bacterial resistance to disinfectants, desiccation, UV exposure, and other stresses, reducing the effectiveness of conventional sanitization methods and contributing to persistent contamination (Shah et al., 2025). The eggshell cuticle is the first natural barrier against microbial invasion, reducing pore accessibility and bacterial penetration. However, its protective function declines due to abrasion, washing, storage, and improper sanitization. Damage to the cuticle enhances bacterial attachment and migration, highlighting the importance of preserving cuticle integrity in alternative biological sanitation strategies (Harris et al., 2022). The shell matrix itself provides partial mechanical resistance against bacterial migration through its complex calcium

carbonate architecture. Nevertheless, microscopic cracks, structural irregularities, and enlarged pore openings substantially reduce this protective capacity (Supaphol, 2026). These contamination mechanisms also explain why simple aqueous washing frequently provides limited microbial reduction. Water alone cannot efficiently disrupt firmly established bacterial adhesion or biofilm matrices and may even facilitate bacterial movement through shell pores if inappropriate washing temperatures are used (Gole et al., 2014; Samiullah et al., 2013). Consequently, modern biological sanitation strategies increasingly incorporate compounds capable of simultaneously damaging bacterial membranes, inhibiting biofilm formation, and maintaining prolonged antimicrobial activity on the shell surface (Melo et al., 2019).




Figure 1. Mechanisms of action of natural antimicrobials against *Salmonella*.

Natural plant based antimicrobial agents

Essential oils

The antimicrobial efficacy of EOs primarily depends on their hydrophobic nature, which facilitates

penetration through the outer membrane of Gram-negative bacteria such as *Salmonella* (Dai et al., 2026). After diffusing into the phospholipid bilayer, bioactive compounds disrupt membrane integrity, increase permeability, collapse the proton motive force, interfere with ATP synthesis, and promote leakage of intracellular constituents including

nucleic acids, proteins, and ions. At higher concentrations, irreversible membrane damage ultimately results in bacterial cell death (**Fig. 1**) (Guimarães et al., 2019). Besides membrane disruption, several EOs constituents also interfere with quorum sensing pathways and inhibit biofilm formation, thereby reducing bacterial persistence on eggshell surfaces (Nesa et al., 2025).

Thymol and carvacrol

Among all phytochemicals investigated for eggshell sanitation, thymol and carvacrol are consistently reported as two of the most potent antimicrobial compounds. These structurally related phenolic monoterpenes, predominantly found in thyme and oregano EOs, exhibit exceptional antimicrobial activity because their hydroxyl groups interact strongly with bacterial membrane phospholipids, resulting in severe membrane destabilization and rapid cytoplasmic leakage (Guimarães et al., 2019). Experimental evidence demonstrates that carvacrol rapidly inactivates *S. Enteritidis* on contaminated shell eggs. In one of the earliest postharvest washing studies, treatments containing carvacrol at concentrations of 0.5% and 0.75% completely eliminated detectable *S. Enteritidis*, even when eggshells were artificially contaminated with 3% chicken fecal material, outperforming conventional chlorine washing under identical experimental conditions (Upadhyaya et al., 2013). Similar observations were reported in subsequent investigations where postharvest washing with 0.25–0.50% carvacrol reduced *S. Enteritidis* to undetectable levels within only 30 seconds, highlighting the rapid antimicrobial kinetics of this phytochemical (Upadhyaya et al., 2014). The effectiveness of thymol has likewise been demonstrated in edible coating systems. Sweet potato starch coatings incorporating thyme EOs at concentrations of 2%, 4%, and 6% significantly inhibited *S. enterica* during storage (Sharaf Eddin & Tahergorabi, 2019). Although the 6% formulation maintained bacterial populations below detectable levels throughout storage, the 4% concentration was identified as the optimum formulation because it provided strong antimicrobial activity while maintaining desirable coating characteristics and

overall egg quality. Beyond direct bactericidal activity, thymol also contributes to biofilm inhibition. Recent evidence indicates that thymol combined with bacteriophage CAU-SEP-3 synergistically disrupted *S. Enteritidis* biofilms on quail eggs without negatively affecting egg quality. Such synergistic approaches are particularly attractive because they permit lower concentrations of each antimicrobial agent while simultaneously improving efficacy against mature biofilms that are otherwise highly resistant to conventional disinfectants (Han et al., 2024).

Trans-cinnamaldehyde

Trans-cinnamaldehyde, the principal aromatic aldehyde of cinnamon EO, has emerged as one of the most extensively investigated phytochemicals for eggshell decontamination. Its antimicrobial activity originates from disruption of membrane-associated proteins, inhibition of energy metabolism, alteration of intracellular pH homeostasis, and suppression of biofilm-associated gene expression, making it highly effective against *Salmonella* under both planktonic and sessile conditions (Shah et al., 2025). Several studies have demonstrated remarkable antimicrobial performance of cinnamaldehyde-based treatments. Washing shell eggs with trans-cinnamaldehyde at concentrations between 0.25% and 0.75% rapidly eliminated *S. Enteritidis*, while fumigation with 1% trans-cinnamaldehyde for 20 min reduced *Salmonella* to undetectable levels on embryonated eggshells and prevented bacterial recovery until day 18 of incubation (Upadhyaya et al., 2015). These findings indicate that gaseous delivery systems may provide highly uniform antimicrobial coverage across complex eggshell surfaces while minimizing excessive liquid exposure. The introduction of nanoemulsion technology has further enhanced cinnamaldehyde performance. Allen et al. (2023) reported that trans-cinnamaldehyde nanoemulsions containing only 0.06–0.48% active compound achieved approximately 2–2.5 log₁₀ CFU/egg reductions after only 1 min of washing without altering eggshell color. Such improvements largely arise from the reduced droplet size of nanoemulsions, which increases surface contact area, enhances bacterial

membrane interactions, and improves aqueous dispersion of this highly hydrophobic compound.

Eugenol

Eugenol, the principal phenolic constituent of clove oil, represents another highly effective antimicrobial compound for eggshell sanitation. Similar to carvacrol, eugenol destabilizes bacterial membranes by disrupting phospholipid organization and increasing membrane permeability. Additionally, its phenolic hydroxyl group contributes to oxidative stress induction and protein denaturation, enhancing bactericidal activity against Gram-negative pathogens (Guimarães et al., 2019). Postharvest washing experiments consistently demonstrate excellent efficacy of eugenol against *Salmonella*. Concentrations of 0.5% and 0.75% completely eliminated detectable *S. Enteritidis* from contaminated shell eggs, including under conditions simulating heavy organic contamination with chicken droppings (Upadhyaya et al., 2013). Likewise, eugenol treatments at concentrations as low as 0.25–0.50% reduced *Salmonella* below detectable limits within 30 seconds of washing (Upadhyaya, 2014). More recently, vapor-phase delivery systems have significantly expanded the practical applications of eugenol. Incorporation of clove oil vapor into antimicrobial paper egg trays at approximately 40 µg/g, followed by UV-C irradiation, extended egg shelf life by approximately five-fold while exhibiting potent antibacterial activity against *Salmonella*, *E. coli*, and *S. aureus*. These findings demonstrate that continuous vapor release during storage may provide long-term antimicrobial protection beyond the immediate effects achieved by conventional washing treatments (Phothisuwan et al., 2020).

Limonene and other citrus EOs

Citrus-derived EOs have recently attracted considerable attention because of their pleasant aroma, low toxicity, and compatibility with active packaging systems (Bertolo et al., 2025). Among them, limonene exhibits broad antimicrobial activity by disrupting bacterial membranes and altering membrane-associated enzyme systems while offering superior consumer acceptance compared

with some strongly aromatic phenolic compounds. Cold limonene vapor generated from nanoemulsions has demonstrated particularly impressive antimicrobial performance. Nanoemulsion concentrations of 0.04%, 0.08%, and 0.12% were evaluated, with 0.08% representing the minimum effective concentration when combined with ultraviolet treatment (Matan & Khunjan, 2026). Under these conditions, bacterial populations of *Salmonella* Enteritidis and *Staphylococcus aureus* decreased by approximately 6–7 log₁₀, indicating nearly complete microbial elimination while preserving egg quality. Comparable success has been reported for citronella EOs. Vaporized citronella nanoemulsions applied at 40, 80, and 120 µL/mL generated thin antimicrobial films on quail eggs, with 80 µL/mL applied for 10 min providing the optimal balance between antimicrobial efficacy and coating uniformity while extending storage life (Kabploy et al., 2025). Similarly, orange oil incorporated into antimicrobial paper egg trays at concentrations between 0.001% and 0.004% (w/w) effectively suppressed pathogenic bacteria for up to 14 days, illustrating the considerable potential of volatile EOs in active packaging systems (Krittika Kabploy et al., 2023).

Plant and fruit extracts

Unlike purified EOs constituents, plant extracts contain diverse mixtures of phenolic acids, flavonoids, tannins, alkaloids, organic acids, and other secondary metabolites that collectively exert antimicrobial and antioxidant activities. The simultaneous presence of multiple bioactive compounds frequently generates synergistic interactions, enabling broader antimicrobial spectra while reducing the concentrations required for effective bacterial inhibition. Moreover, many plant extracts exhibit lower volatility than EOs, making them particularly suitable for incorporation into edible coatings intended for prolonged storage (Teshome et al., 2022).

Green tea (*Camellia sinensis*) is among the most extensively investigated botanical extracts for egg preservation because of its exceptionally high concentration of catechins, particularly epigallocatechin gallate, which possesses potent

antimicrobial and antioxidant properties. These polyphenols interact with bacterial membranes, increase membrane permeability, inhibit essential enzymes, and induce oxidative stress, ultimately resulting in bacterial death (Friedman et al., 2006). Songsamoe et al. (2022) evaluated combinations of green tea extract and peppermint EOs for sanitizing eggs stored at ambient temperature. Among all formulations tested, a combination containing 10% green tea extract and 0.15% peppermint oil achieved complete antibacterial activity against the tested pathogenic bacteria, extending egg shelf life from approximately 7 days to at least 14 days under room temperature. The study further proposed spray application as a practical industrial approach because it provided uniform antimicrobial coverage while minimizing mechanical damage to the eggshell.

Pomegranate peel represents an abundant source of ellagic acid, tannins, and other phenolic antioxidants capable of suppressing microbial growth while simultaneously delaying oxidative deterioration. Thiangtae et al. (2026) developed gelatin-based coatings containing 1.0% and 1.5% pomegranate peel extract, demonstrating effective preservation of washed eggs during 28 days of tropical storage. The improved preservation resulted from complementary mechanisms whereby the gelatin matrix reduced moisture transfer while ellagic acid-rich extract minimized oxidative degradation and inhibited microbial proliferation. This dual protective mechanism illustrates how fruit-derived polyphenols can simultaneously improve microbial safety and physicochemical stability.

Garlic (*Allium sativum*) has long been recognized for its antimicrobial properties owing primarily to sulfur-containing compounds such as allicin and ajoene (Razavi Rohani et al., 2011). Incorporation of garlic extract into polyvinyl alcohol/chitosan/montmorillonite nanocomposite coatings significantly improved egg preservation during four weeks of storage. The formulation containing 4% garlic extract exhibited superior maintenance of shell strength and internal egg quality while displaying concentration-dependent antimicrobial activity, particularly against Gram-positive bacteria (Yousefi Zirabi & khademi

shurmasti, 2024). Recent studies have increasingly explored novel botanical materials possessing both antimicrobial and antioxidant activities. Bioactive coatings enriched with *Cordia verbenacea* aqueous extract, for example, were recently shown to preserve eggs under non-refrigerated conditions for 25 days. Remarkably, the coating containing 100% plant extract completely eliminated detectable microbial contamination on eggshells, reducing microbial counts from 70.0 ± 5.8 CFU/egg in untreated controls to zero, highlighting the substantial potential of plant-derived coatings for long-term microbial control (Soares et al., 2026). Cinnamon hydrosol, a by-product of EOs extraction, has been successfully incorporated into pectin coatings to improve coating homogeneity and prolong egg shelf life despite containing relatively low EOs concentrations (< 1 g/L) (Didar, 2019).

Application methods on eggshells

Current application strategies include immersion (dipping), spraying, edible coatings, fumigation, vapor-phase treatments, nanoemulsion-based delivery, and incorporation into active packaging materials, each possessing distinct advantages and limitations (**Table 1**) depending on the processing environment and intended storage period (Moradi et al., 2021; Oliveira et al., 2022).

Immersion and washing treatments

Immersion is one of the simplest and most widely investigated approaches for applying natural antimicrobial compounds to eggshells. In this technique, eggs are submerged in antimicrobial solutions for a predetermined period, allowing direct contact between the antimicrobial agent and bacterial cells attached to the shell surface. Because immersion is compatible with existing commercial egg-washing equipment, it represents one of the most practical strategies for industrial implementation. Upadhyaya et al. (2013) evaluated postharvest washing solutions containing carvacrol, eugenol, β -resorcylic acid, and trans-cinnamaldehyde against *S. Enteritidis* artificially inoculated onto shell eggs. Washing with 0.5% and 0.75% carvacrol or eugenol completely eliminated detectable *S. Enteritidis*, even when eggs were

contaminated with 3% chicken fecal material, demonstrating excellent efficacy under conditions simulating commercial contamination. Similar reductions were obtained using trans-cinnamaldehyde, indicating that immersion can rapidly inactivate surface-associated bacteria when suitable phytochemicals are employed. Subsequent investigations further demonstrated the rapid antimicrobial kinetics achievable through immersion. Immersion treatments present several operational challenges. Improper wash-water

temperature may facilitate bacterial penetration through eggshell pores owing to pressure differentials generated during cooling. Furthermore, repeated use of washing solutions may increase the risk of cross-contamination if antimicrobial concentrations decline below effective levels (El-Sabrou et al., 2026; Oliveira et al., 2022). Consequently, strict control of solution composition, temperature, and replacement frequency remains essential for maintaining microbiological safety.

Table 1. Comparison of delivery systems for natural antimicrobials applied to shell eggs

Delivery system	Representative antimicrobial	Main advantages	Main limitations	Industrial potential	References
Washing/Immersion	Carvacrol, Eugenol, Cinnamaldehyde	Rapid microbial reduction; simple process	Possible cross-contamination; limited residual activity	High	(Upadhyaya et al., 2014; Upadhyaya et al., 2013)
Spray application	Green tea extract + peppermint oil	Uniform coverage; low solution consumption; easy automation	Coverage depends on nozzle design	Very high	(Songsamoe et al., 2022)
Edible coating	Chitosan, alginate, pectin, whey protein	Sustained antimicrobial release; improved egg quality	Additional drying step required	Very high	(Jin et al., 2013; Supaphol, 2026)
Nanoemulsion	Trans-cinnamaldehyde; limonene	Enhanced stability; improved solubility; lower effective concentration	Higher production cost	High	(Allen et al., 2023; Matan & Khunjan, 2026)
Vapor fumigation	Trans-cinnamaldehyde	Uniform shell coverage; suitable for hatching eggs	Specialized equipment required	Moderate–High	(Upadhyaya et al., 2015)
Active packaging	Orange oil; clove oil; <i>Litsea cubeba</i> oil	Continuous antimicrobial release during storage	Packaging development costs	Very high	(Kabploy et al., 2023; Phothisuwan et al., 2020)
Nanocomposite coating	Garlic extract; dill oil	Improved mechanical strength and controlled release	More complex formulation	Emerging	(Yousefi Zirabi & khademi shurmasti, 2024)
Hurdle technology	EOs + bacteriophage/UV/pulse electric field	Synergistic antimicrobial effect; lower antimicrobial doses	Process optimization required	Future commercial application	(Han et al., 2024; Matan & Khunjan, 2026).

Spray application

Recent studies indicate that spray application may achieve antimicrobial efficacy comparable to immersion while improving process sustainability. Songsamoe et al. (2022) proposed spray application of a formulation containing 10% green tea extract

and 0.15% peppermint EO, which successfully extended shelf life from approximately 7 to 14 days under ambient storage. The authors emphasized that spraying ensured homogeneous surface coverage while minimizing mechanical abrasion of the eggshell cuticle, an important advantage for preserving the egg's natural protective barrier.

Compared with immersion, spray application also allows easier adjustment of antimicrobial dosage according to contamination level and production speed. Spraying minimizes the accumulation of contaminated wash water and significantly reduces opportunities for cross-contamination between eggs. Nevertheless, adequate nozzle design, droplet size optimization, and uniform spray distribution remain critical engineering parameters influencing antimicrobial performance in industrial environments.

Edible coating application

Edible coatings represent one of the most sophisticated antimicrobial delivery systems currently available for shell eggs. Rather than providing only transient microbial reduction immediately after treatment, coating technologies establish a semipermeable protective layer that remains attached to the eggshell throughout storage. This layer gradually releases antimicrobial compounds while simultaneously reducing moisture loss, limiting gas exchange, and preserving internal egg quality (Ezazi et al., 2021). Numerous studies have demonstrated that coating systems substantially outperform direct antimicrobial application during prolonged storage. Chitosan coatings incorporating 1.0% lauric arginate ester reduced *Salmonella* populations by approximately $5.2 \log_{10}$ CFU/cm² (Jin et al., 2013), whereas pectin-based coatings containing carvacrol, eugenol, and 0.75% β -resorcylic acid completely eliminated detectable *S. Enteritidis* after only three days of refrigerated storage (Upadhyaya et al., 2016). These findings indicate that sustained antimicrobial release rather than immediate bacterial killing is largely responsible for the superior long-term performance of coating technologies. An additional advantage of edible coatings is their multifunctionality. Besides serving as antimicrobial reservoirs, coatings improve Haugh unit values, reduce albumen thinning, minimize weight loss, and maintain eggshell strength during storage. Consequently, edible coatings simultaneously enhance microbial safety and commercial egg quality, providing significant advantages over conventional

disinfectants that primarily target microorganisms alone (Avramescu et al., 2020; Supaphol, 2026).

Nanoemulsion-based delivery systems

The hydrophobic nature of most EOs considerably limits their direct application in aqueous sanitizing solutions. Nanoemulsion technology addresses this challenge by dispersing EOs droplets into nanoscale particles, thereby improving water dispersibility, physical stability, surface contact area, and controlled antimicrobial release (Ghodsi Sheykhjan et al., 2025). Allen et al. (2023) demonstrated that nanoemulsions containing only 0.06–0.48% trans-cinnamaldehyde achieved approximately 2–2.5 $\log_{10}$ CFU/egg reductions after merely one minute of washing, without causing measurable alterations in eggshell color. Compared with conventional emulsions, the nano-scale droplets exhibited improved interaction with bacterial membranes, leading to more efficient microbial inactivation despite substantially lower active ingredient concentrations. Nanoemulsion technology has also proven highly effective for vapor-based treatments. Matan and Khunjan (2026) reported that cold limonene nanoemulsion vapor at 0.08%, combined with ultraviolet treatment, reduced *S. Enteritidis* and *S. aureus* populations by approximately 6–7 $\log_{10}$, while preserving egg quality. Similarly, citronella nanoemulsion vapor applied at 80 μ L/mL for 10 min generated a uniform antimicrobial coating that effectively extended quail egg shelf life (Kabploy et al., 2025; Matan & Khunjan, 2026).

Vapor-phase fumigation and active packaging

Unlike liquid-based application methods, vapor-phase technologies deliver volatile antimicrobial compounds through the surrounding atmosphere rather than by direct surface contact. This approach provides highly uniform antimicrobial exposure over complex shell geometries while minimizing excessive moisture accumulation. Upadhyaya et al. (2015) demonstrated that fumigation with 1% trans-cinnamaldehyde for 20 min completely eliminated detectable *S. Enteritidis* from embryonated eggshells, with no bacterial recovery observed during incubation. These findings suggest that gaseous antimicrobial delivery may be particularly

suitable for hatching eggs where excessive liquid exposure should be avoided. Recent advances have extended this concept to active packaging systems. Antimicrobial paper egg trays containing orange oil, clove oil, or *Litsea cubeba* EOs gradually release volatile antimicrobial compounds throughout storage. Orange oil incorporated at concentrations between 0.001% and 0.004% (w/w) effectively suppressed pathogenic bacteria for up to 14 days, whereas clove oil vapor combined with UV-C irradiation extended egg shelf life approximately five-fold while inhibiting *Salmonella*, *E. coli*, and *S. aureus* (Kabploy et al., 2023; Phothisuwan et al., 2020).

Operational challenges and quality assessment

Sensory impact

One of the principal challenges associated with natural antimicrobial compounds is their potential influence on the sensory characteristics of eggs. EOs possess strong volatile aromas that may alter the odor, flavor, or overall acceptability of eggs when applied at high concentrations. This issue is particularly important because consumer purchasing decisions are strongly influenced by external appearance and perceived freshness, even before the product is consumed (Yazdanbakhsh et al., 2026). The intensity of sensory alterations largely depends on the chemical composition and concentration of the antimicrobial compound. Phenolic-rich EOs such as oregano, thyme, clove, and cinnamon generally exhibit superior antimicrobial activity but also possess stronger characteristic aromas than citrus-derived EOs (Falleh, 2025; Hyldgaard et al., 2012). Consequently, identifying the minimum effective antimicrobial concentration has become a major objective of formulation studies. Several investigations included in this review demonstrated that appropriate formulation strategies can effectively minimize undesirable sensory changes. Allen et al. (2023) reported that washing shell eggs with 0.06–0.48% trans-cinnamaldehyde nanoemulsion produced approximately 2–2.5 log₁₀ CFU/egg reductions of *S.*

Enteritidis without causing measurable changes in eggshell color. The nanoemulsion system improved antimicrobial efficiency while permitting substantially lower concentrations of cinnamaldehyde than conventional emulsions, thereby reducing the likelihood of undesirable sensory effects. Similarly, Sharaf Eddin and Tahergorabi (2019) observed that sweet potato starch coatings containing 4% thyme EO provided the best overall balance between antimicrobial efficacy and coating quality. Although the 6% formulation exhibited slightly greater antimicrobial activity, increasing EOs concentration adversely affected coating properties, illustrating the importance of optimizing formulations to simultaneously satisfy microbiological and sensory requirements.

Stability of bioactive compounds during storage

Another important limitation concerns the chemical stability of natural antimicrobial compounds throughout storage and distribution. Many EOs constituents are highly volatile and susceptible to oxidation, photodegradation, and thermal degradation, leading to progressive reductions in antimicrobial activity during prolonged storage. This instability represents one of the principal reasons why direct application of EOs frequently provides only short-term protection (Ganosi et al., 2023; Singh & Pulikkal, 2022). Active packaging systems provide an additional strategy for maintaining antimicrobial activity during commercial distribution. Rather than releasing the entire antimicrobial dose immediately after application, packaging materials containing volatile EOs gradually emit antimicrobial vapors throughout storage. For example, antimicrobial paper trays containing orange oil effectively inhibited pathogenic bacteria for 14 days (Kabploy et al., 2023), whereas clove oil incorporated into paper trays extended egg shelf life by approximately five-fold when combined with UV-C treatment (Phothisuwan et al., 2020). These findings indicate that controlled-release technologies substantially improve the long-term stability and effectiveness of volatile phytochemicals.

Conclusion and future perspectives

Bioactive compounds such as thymol, carvacrol, eugenol, cinnamaldehyde, limonene, and diverse plant phenolics effectively inhibit *Salmonella* by disrupting bacterial membranes, interfering with cellular metabolism, and suppressing biofilm formation. Numerous studies have demonstrated that these natural compounds can significantly reduce bacterial contamination while maintaining egg quality and extending shelf life. Recent advances in nanoemulsion technology and active packaging have further improved the stability and antimicrobial performance of EOs, increasing their suitability for commercial application. Despite remaining challenges related to formulation standardization, regulatory approval, and industrial scalability, the available evidence strongly supports the future integration of EOs and plant extracts into sustainable egg sanitation strategies. Continued research focusing on optimized delivery systems and commercial validation will accelerate their adoption as environmentally friendly alternatives to conventional chemical disinfectants.

Conflicts of interest

The authors declare no conflicts of interest.

Disclaimer

Author hereby declares that NO generative AI technologies, have been used during manuscript preparation.

References

- Alkan, D., & Yemenicioğlu, A. (2016). Potential application of natural phenolic antimicrobials and edible film technology against bacterial plant pathogens. *Food Hydrocolloids*, 55, 1–10. <https://doi.org/10.1016/j.foodhyd.2015.10.025>
- Allen, J., Balasubramanian, B., Rankin, K., Shah, T., Donoghue, A. M., Upadhyaya, I., Sartini, B., Luo, Y., & Upadhyay, A. (2023). Trans-cinnamaldehyde nanoemulsion wash inactivates *Salmonella* Enteritidis on shelled eggs without affecting egg color. *Poultry Science*, 102(4), 102523. <https://doi.org/10.1016/j.psj.2023.102523>
- Avramescu, S. M., Butean, C., Popa, C. V., Ortan, A., Moraru, I., & Temocico, G. (2020). Edible and functionalized films/coatings—performances and perspectives. *Coatings*, 10(7), 687. <https://doi.org/10.3390/coatings10070687>

Bertolo, M. R. V., Pereira, T. S., dos Santos, F. V., Facure, M. H. M., dos Santos, F., Teodoro, K. B. R., Mercante, L. A., & Correa, D. S. (2025). Citrus wastes as sustainable materials for active and intelligent food packaging: Current advances. *Comprehensive Reviews in Food Science and Food Safety*, 24(2), e70144. <https://doi.org/https://doi.org/10.1111/1541-4337.70144>

Dai, J., Xue, S., Bai, M., Chiu, T., Aziz, T., Al-Asmari, F., Jalal, R. S., Cui, H., & Lin, L. (2026). Structure-activity relationships governing the membrane targeted antimicrobial mechanisms of essential oils for food pathogen control. *Food Bioscience*, 82, 109422. <https://doi.org/10.1016/j.fbio.2026.109422>

Didar, Z. (2019). Effects of coatings with pectin and *Cinnamomum verum* hydrosol included pectin on physical characteristics and shelf life of chicken eggs stored at 30°C. *Nutrition and Food Sciences Research*, 6(4), 39–45.

El-kashef, M. M. A., & El Sabry, M. I. (2025). Egg sanitation with ginger and garlic solutions affects embryonic development, hatchability, blood parameters and post-hatch performance of Japanese quail. *Tropical Animal Health and Production*, 57(7), 338. <https://doi.org/10.1007/s11250-025-04556-8>

El-Sabrou, K., Elahi, U., Ahmad, S., Vallone, L., & Buonaiuto, G. (2026). Sustainable biosecurity: harnessing natural sanitisers for hatching egg sanitation. *World's Poultry Science Journal*, 82(2), 651–679. <https://doi.org/10.1080/00439339.2026.2628865>

Elkomy, A. E., El-Saadany, A. S., Shreif, E. Y., Bayoumi, A. A., Abd El-Maged, M. H., Alagawany, M., Saleh, A. A., Cho, S., Kim, I. H., & Eltahan, H. M. (2023). Anise and grape seed oils as a feed additive to improve the performance, immune response, and antioxidant activity and reduce caecal pathogenic microbes of quail. *Archives Animal Breeding*, 66(4), 379–390. <https://doi.org/10.5194/aab-66-379-2023>

Ezazi, A., Javadi, A., Jafarizadeh-Malmiri, H., & Mirzaei, H. (2021). Development of a chitosan-propolis extract edible coating formulation based on physico-chemical attributes of hens' eggs: Optimization and characteristics edible coating of egg using chitosan and propolis. *Food Bioscience*, 40, 100894. <https://doi.org/10.1016/j.fbio.2021.100894>

Falleh, H. (2025). Demystifying the power of essential oils: a review of their antibacterial properties and potential as natural food preservatives. *EXCLI Journal*, 24, 828–850. <https://doi.org/10.17179/excli2025-8439>

Friedman, M., Henika, P. R., Levin, C. E., Mandrell, R. E., & Kozukue, N. (2006). Antimicrobial activities of tea catechins and theaflavins and tea extracts against *Bacillus cereus*. *Journal of Food Protection*, 69(2), 354–361. <https://doi.org/10.4315/0362-028X-69.2.354>

Ganosi, E., Barda, C., Grafakou, M.-E., Rallis, M. C., & Skaltsa, H. (2023). An in-depth stability study of the essential oils from *Mentha × piperita*, *Mentha spicata*, *Origanum vulgare*, and *Thymus vulgaris*: The impact of thermal and storage conditions. *Separations*, 10(9), 488. <https://doi.org/10.3390/separations10090488>

- Ghodsi Sheykhjan, M., Fazlara, A., Hojjati, M., & Alizadeh Behbahani, B. (2025). Active food packaging strategies: Promoting mutton quality using *Lepidium sativum* coatings and nanoemulsions of *Dracocephalum moldavica* essential oil. *Applied Food Research*, 5(1), 101021. <https://doi.org/10.1016/j.afres.2025.101021>
- Gole, V. C., Chousalkar, K. K., Roberts, J. R., Sexton, M., May, D., Tan, J., & Kiermeier, A. (2014). Effect of egg washing and correlation between eggshell characteristics and egg penetration by various *Salmonella* Typhimurium strains. *Plos One*, 9(3), e90987. [10.1371/journal.pone.0090987](https://doi.org/10.1371/journal.pone.0090987)
- Guimarães, A. C., Meireles, L. M., Lemos, M. F., Guimarães, M. C., Endringer, D. C., Fronza, M., & Scherer, R. (2019). Antibacterial activity of terpenes and terpenoids present in essential oils. *Molecules*, 24(13), 2471. <https://doi.org/10.3390/molecules24132471>
- Han, S., Choi, M. W., Byun, K.-H., Kim, B. H., Song, M. S., Kang, I., & Ha, S.-D. (2024). Characterization of *Salmonella* ser. Enteritidis-specific bacteriophages and biocontrol strategy to reduce *S. Enteritidis* on egg products using bacteriophages and essential oil compounds. *Food Control*, 160, 110304. <https://doi.org/10.1016/j.foodcont.2024.110304>
- Harris, C. E., Josselson, L. N. B., Bourassa, D. V., & Buhr, R. J. (2022). Examination of the impact of eggshell cuticle and membranes on *Salmonella* Enteritidis or Typhimurium recovery from inoculated and stored eggs. *Journal of Applied Poultry Research*, 31(4), 100297. <https://doi.org/10.1016/j.japr.2022.100297>
- Hylgaard, M., Mygind, T., & Meyer, R. L. (2012). Essential oils in food preservation: mode of action, synergies, and interactions with food matrix components. *Frontiers in Microbiology*, 3, 12. <https://doi.org/10.3389/fmicb.2012.00012>
- Illera, A. E., Souza, V. R., Nikmaram, N., Tang, L., & Keener, K. M. (2022). High voltage atmospheric cold plasma decontamination of *Salmonella* enteritidis on chicken eggs. *Innovative Food Science & Emerging Technologies*, 82, 103210. <https://doi.org/10.1016/j.ifset.2022.103210>
- Jin, T. Z., Gurtler, J. B., & Li, S.-Q. (2013). Development of antimicrobial coatings for improving the microbiological safety and quality of shell eggs. *Journal of Food Protection*, 76(5), 779–785. <https://doi.org/10.4315/0362-028X.JFP-12-460>
- Kabpoy, K., Khunjan, K., & Matan, N. (2025). Underlying mechanisms of citronella oil vapor generated by an ultrasonic humidifier in inhibiting pathogenic bacteria and extending the shelf life of quail eggs. *ES Food and Agroforestry*, 19. <https://doi.org/10.30919/faf1400>
- Kabpoy, Krittika, Sukmas, S., Phothisuwan, S., Sinthupachee, A., Kongchoosi, N., & Matan, N. (2023). Development of an antimicrobial fungal egg tray containing orange oil and smoke for eggs preservation at room temperature. *Poultry Science*, 102(9), 102885. <https://doi.org/10.1016/j.psj.2023.102885>
- Karnwal, A., & Malik, T. (2024). Exploring the untapped potential of naturally occurring antimicrobial compounds: novel advancements in food preservation for enhanced safety and sustainability. *Frontiers in Sustainable Food Systems*, 8, 1307210. <https://doi.org/10.3389/fsufs.2024.1307210>
- Konfo, T. R. C., Djouhou, F. M. C., Koudoro, Y. A., Dahouenon-Ahoussi, E., Avlessi, F., Sohounhloue, C. K. D., & Simal-Gandara, J. (2023). Essential oils as natural antioxidants for the control of food preservation. *Food Chemistry Advances*, 2, 100312. <https://doi.org/10.1016/j.focha.2023.100312>
- Matan, N., & Khunjan, K. (2026). Cold limonene vapor and UV treatment against *Salmonella* Enteritidis and *Staphylococcus aureus* in quail eggs: Potential mode of action. *Foodborne Pathogens and Disease*, 23(6), 384–390. <https://doi.org/10.1089/fpd.2025.0035>
- Melo, E. F., Clímaco, W. L. S., Triginelli, M. V., Vaz, D. P., de Souza, M. R., Baião, N. C., Pompeu, M. A., & Lara, L. J. C. (2019). An evaluation of alternative methods for sanitizing hatching eggs. *Poultry Science*, 98(6), 2466–2473. <https://doi.org/10.3382/ps/pez022>
- Moradi, M., Kousheh, S. A., Razavi, R., Rasouli, Y., Ghorbani, M., Divsalar, E., Tajik, H., Guimarães, J. T., & Ibrahim, S. A. (2021). Review of microbiological methods for testing protein and carbohydrate-based antimicrobial food packaging. *Trends in Food Science & Technology*, 111, 595–609. <https://doi.org/10.1016/j.tifs.2021.03.007>
- Nesa, S. R., Mizan, M. F. R., Meghla, N. S., Kang, I., & Ha, S.-D. (2025). Exploring the effects of phenolic compounds and essential oils in poultry: A sustainable strategy to combat *Salmonella* biofilm infections. *Poultry Science*, 104(12), 106056. <https://doi.org/10.1016/j.psj.2025.106056>
- Oliveira, G. D., McManus, C., Salgado, C. B., & dos Santos, V. M. (2022). Effects of sanitizers on microbiological control of hatching eggshells and poultry health during embryogenesis and early stages after hatching in the last decade. *Animals*, 12(20), 2826. <https://doi.org/10.3390/ani12202826>
- Phothisuwan, S., Preechatiwong, W., & Matan, N. (2020). Enhancement of antibacterial activity of essential oil vapor released from a paper egg tray in combination with UV-C radiation against pathogenic bacteria on chicken eggs. *Journal of Food Processing and Preservation*, 44(10), e14794. <https://doi.org/https://doi.org/10.1111/jfpp.14794>
- Raikwar, G., Kumar, D., Mohan, S., & Dahiya, P. (2024). Synergistic potential of essential oils with antibiotics for antimicrobial resistance with emphasis on mechanism of action: A review. *Biocatalysis and Agricultural Biotechnology*, 61, 103384. <https://doi.org/10.1016/j.bcab.2024.103384>
- Razavi Rohani, S. M., Moradi, M., Mehdizadeh, T., Saei-Dehkordi, S. S., & Griffiths, M. W. (2011). The effect of nisin and garlic (*Allium sativum* L.) essential oil separately and in combination on the growth of *Listeria monocytogenes*. *LWT - Food Science and Technology*, 44(10), 2260–2265. <https://doi.org/10.1016/j.lwt.2011.07.020>

- Samiullah, Chousalkar, K. K., Roberts, J. R., Sexton, M., May, D., & Kiermeier, A. (2013). Effects of egg shell quality and washing on *Salmonella* Infantis penetration. *International Journal of Food Microbiology*, 165(2), 77–83. <https://doi.org/10.1016/j.ijfoodmicro.2013.05.002>
- Shah, T., Zhu, C., Shah, C., Upadhyaya, I., & Upadhyay, A. (2025). Trans-cinnamaldehyde nanoemulsion reduces *Salmonella* Enteritidis biofilm on steel and plastic surfaces and downregulates expression of biofilm associated genes. *Poultry Science*, 104(5), 105086. <https://doi.org/10.1016/j.psj.2025.105086>
- Sharaf Eddin, A., & Tahergorabi, R. (2019). Efficacy of sweet potato starch-based coating to improve quality and safety of hen eggs during storage. *Coatings*, 9(3), 205. <https://doi.org/10.3390/coatings9030205>
- Singh, I. R., & Pulikkal, A. K. (2022). Preparation, stability and biological activity of essential oil-based nano emulsions: A comprehensive review. *OpenNano*, 8, 100066. <https://doi.org/10.1016/j.onano.2022.100066>
- Soares, J. G., Serafini, S., Picoli, F., Araújo, D. N., Boiago, M. M., Galvão, A. C., & Robazza, W. D. (2026). Citric Pectin–*Cordia verbenacea* bioactive coatings to preserve egg quality under non-refrigerated conditions using machine learning approaches. *Foods*, 15(5), 879. <https://doi.org/10.3390/foods15050879>
- Songsamoe, S., Kabploy, K., Khunjan, K., & Matan, N. (2022). The combined effect of green tea and peppermint oil against pathogenic bacteria to extend the shelf life of eggs at ambient temperature and the mode of action. *Journal of Food Safety*, 42(1), e12945. <https://doi.org/https://doi.org/10.1111/jfs.12945>
- Supaphol, P. (2026). Food coatings for egg preservation: a comprehensive review of materials, mechanisms and performance. *World's Poultry Science Journal*, 1–42. <https://doi.org/10.1080/00439339.2026.2663103>
- Teshome, E., Forsido, S. F., Rupasinghe, H. P. V., & Olika Keyata, E. (2022). Potentials of natural preservatives to enhance food safety and shelf Life: A review. *The Scientific World Journal*, 2022(1), 9901018. <https://doi.org/https://doi.org/10.1155/2022/9901018>
- Thiangtae, S., Srikijkasemwat, K., Kaewhom, P., Kongrith, C., Polyorach, S., Klompanya, A., & Thammakarn, C. (2026). Enhancing the shelf-life of washed eggs using a gelatin-based coating enriched with pomegranate peel extract under tropical storage conditions. *Applied Food Research*, 6(1), 101946. <https://doi.org/10.1016/j.afres.2026.101946>
- Upadhyaya, I., Kollanoor-Johny, A., Darre, M. J., & Venkitanarayanan, K. (2014). Efficacy of plant-derived antimicrobials for reducing egg-borne transmission of *Salmonella* Enteritidis. *Journal of Applied Poultry Research*, 23(2), 330–339. <https://doi.org/10.3382/japr.2014-00941>
- Upadhyaya, Indu, Upadhyay, A., Kollanoor-Johny, A., Baskaran, S. A., Mooyottu, S., Darre, M. J., & Venkitanarayanan, K. (2013). Rapid inactivation of *Salmonella* Enteritidis on shell eggs by plant-derived antimicrobials. *Poultry Science*, 92(12), 3228–3235. <https://doi.org/10.3382/ps.2013-03126>
- Upadhyaya, Indu, Yin, H.-B., Nair, M. S., Chen, C.-H., Upadhyay, A., Darre, M. J., & Venkitanarayanan, K. (2015). Efficacy of fumigation with Trans-cinnamaldehyde and eugenol in reducing *Salmonella enterica* serovar Enteritidis on embryonated egg shells. *Poultry Science*, 94(7), 1685–1690. <https://doi.org/10.3382/ps/pev126>
- Upadhyaya, Indu, Yin, H.-B., Surendran Nair, M., Chen, C.-H., Lang, R., Darre, M. J., & Venkitanarayanan, K. (2016). Inactivation of *Salmonella* Enteritidis on shell eggs by coating with phytochemicals. *Poultry Science*, 95(9), 2106–2111. <https://doi.org/10.3382/ps/pew152>
- Yazdanbakhsh, H., Alahmoradi, S., Abdolahi-Fard, P., Moradiyan, M. H., Bahraminejad, M., Moradi, D., Assadpour, E., Azizi-Lalabadi, M., & Jafari, S. M. (2026). Essential oil-loaded electrospun fibers/nanofibers for active food packaging. *Advances in Colloid and Interface Science*, 103990. <https://doi.org/10.1016/j.cis.2026.103990>
- Yousefi Zirabi, E., & khademi shurmasti, D. (2024). Monitoring the physicochemical changes of eggs coated with active nanocomposite incorporating garlic extract (*Allium sativum* L.) during storage. *Journal of Food Science and Technology (Iran)*, 21(153), 1–12. <https://doi.org/10.22034/FSCT.21.153.1>
- Zhang, N., Zhang, Y., Yu, T., Tang, M., Liu, M., Zhong, C., Chen, X., Wang, B., Xu, X., & Qin, S. (2026). Antimicrobial mechanism of *Artemisia argyi* essential oil against Carbapenem-resistant *Escherichia coli* by metabolomics analysis. *Journal of Holistic Integrative Pharmacy*, 7(1), 236–248. <https://doi.org/10.1016/j.jhip.2026.03.003>
- Zhang, Y., Huang, K., Tao, N., & Wang, X. (2026). Recent advancements in modification of essential oils and its application in food packaging. *Trends in Food Science & Technology*, 168, 105462. <https://doi.org/10.1016/j.tifs.2025.105462>